WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ :		(11) International Publication Number:	WO 98/48048
C12Q 1/68	A2	(43) International Publication Date:	29 October 1998 (29.10.98)

PCT/GB98/01138 (21) International Application Number: 20 April 1998 (20.04.98) (22) International Filing Date: (30) Priority Data: 21 April 1997 (21.04.97) GB 9707996.6

(71) Applicant (for all designated States except US): CAMBRIDGE UNIVERSITY TECHNICAL SERVICES LTD. [GB/GB]; The Old Schools, Trinity Lane, Cambridge CB2 1TS (GB).

(72) Inventor; and (75) Inventor/Applicant (for US only): BALASUBRAMANIAN, Shankar [GB/GB]; University of Cambridge, Dept. of Chemistry, Lensfield Road, Cambridge CB2 1EW (GB).


(74) Agent: GILL JENNINGS & EVERY; Broadgate House, 7 Eldon Street, London EC2M 7LH (GB).

(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).

Published

Without international search report and to be republished upon receipt of that report.

(54) Title: DNA MUTATION MAPPING BY MULTIPLE ENERGY TRANSFER INTERACTIONS



(57) Abstract

A method for determining the presence and location of a mismatch in a target sequence using Fluorescence Resonance Energy Transfer (FRET). The method comprises contacting the target sequence with at least three labelled oligonucleotide probes capable of hybridising to the natural sequence, in juxtaposition. The central probe is labelled with a donor fluorophore and the probe either side of this is labelled with a distinct acceptor fluorophore. Hybridisation to the target sequence results in resonance between the donor fluorophore and the acceptor fluorophores. A mismatch present in the target sequence will disrupt hybridisation to that region, resulting in a significant alteration to the resonance signal.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM.	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
ΑT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU.	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ.	Azerbaijan	GB	United Kingdom	MC	Monaco	TĐ	Chad
3A	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
3B	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
E	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
F	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
G	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
IJ	Benin	IE	Ireland	MN	Mongolia	ŲA	Ukraine
R	Brazil	1L	Israel	MR	Mauritania	UG	Uganda
ΙΥ	Belarus	IS	Iceland	MW	Malawi	US	United States of America
A	Canada	IТ	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
:G	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
31	Switzerland	KG	Kyrgyzstan	NO	Norway	zw	Zimbabwe
1	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM .	Cameroon		Republic of Korea	PL	Poland		
N.	China	KR	Republic of Korea	PT	Portugal		
วบ	Cuba	KZ	Kazakstan	RO	Romania		
Z	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
E	Germany	Li	Liechtenstein	SD	Sudan		
K	Denmark	LK	Sri Lanka	SE	Sweden		
EE.	Estonia	LR	Liberia	SG	Singapore		

1

\$1 - -----

DNA MUTATION MAPPING BY MULTIPLE ENERGY TRANSFER INTERACTIONS

Field of Invention

5

10

15

20

25

30

35

This invention relates to DNA mismatch screening, especially using FRET-detected hybridisation.

Background of the Invention

There is a general increase in the exploration of gene sequences and function. This has generated a need for new approaches to nucleic acid analysis.

One of the simplest methods for detecting gene sequences is to make use of the specific hybridisation reaction between the target sequence and a suitable probe.

Recently, the use of Fluorescence Resonance Energy Transfer (FRET) has been applied to the detection of hybridised probes. Tyagi et al., Nature Biotech. (1996) 14:303-308, describe the use of FRET to distinguish between hybridised and unhybridised probes in a homogeneous assay. The probes each comprise a stem-and-loop structure with the stem formed by the annealing of two complementary arm sequences either side of the probe sequence. A fluorescent moiety is attached to one arm and a non-fluorescent quenching moiety is attached to the other arm. Separation of the stem structure occurs on hybridisation of the probe to the complementary target sequence. This separates the two moieties and allows fluorescence to occur.

Summary of Invention

This invention is based on a realisation of the utility of FRET as a technique for mapping point mutations on a single strand of DNA.

According to the present invention, a method for determining the presence of a mismatch in a target sequence, comprises contacting the target sequence with first, second and third oligonucleotides capable of hybridising to the natural sequence, in juxtaposition, wherein the oligonucleotides are respectively labelled with first, second and third markers having first, second and third absorption wavelengths and first, second and third

2

emission wavelengths such that there is resonance between either or each of first and second markers and between second and third markers, and observing the presence or absence of each or either resonance.

5 Description of the Drawings

10

15

20

25

30

35

The accompanying drawings are for the purpose of illustration only. In the drawings:

Fig. 1 is a schematic representation of the interaction of labelled oligonucleotides and target sequence in a screening method embodying the invention;

Fig. 2 is a schematic representation of a more specific embodiment of the invention and represents another arrangement of the labels; and

Fig. 3 illustrates the different emission intensities obtained when point mutations are present in the target sequence, using the arrangement of fluorophores shown in Fig. 2.

Description of Invention

The ability to monitor the interactions between nucleic acids is achieved through the detection of sensitised acceptor emission, due to FRET between a donor fluorophore and two different acceptor fluorophores on associating strands. The absence of any of the components induced by a single base mismatch should cause a detectable loss in sensitised acceptor emission.

The system illustrated in Fig. 1 comprises a target strand and three adjacent complementary oligonucleotides, A, B and C. These are typically each 8-13 base pairs in length. A and B are modified at the 5' end with a fluorescein moiety (F). B and C are modified at the 3' end with two distinct types of acceptor fluorophore (X and Y) having different emission maxima and absorption profiles which overlap with the emission profile of fluorescein.

When no mismatches are present, all three oligonucleotides will be bound to the target strand. Thus, when the system is excited at the absorption wavelength of fluorescein (488 nm), FRET will occur and the fluorescence

3

spectrum will show a decrease in the fluorescein emission relative to each singly-bound probe and appearance of two maxima due to secondary emissions from X and Y. In cases where a mismatch exists somewhere within the target sequence, the region in which the error occurs can be fluorescence analysing the by distinguished simply If the mismatch lies within the binding domain spectrum. of oligonucleotide A, no duplex will form in this region and the secondary emission from X will be lost. Similarly, for mismatches within the B region, the signals from X and Y will disappear, and for errors in region C, the Y emission will be absent.

10

15

20

25

30

35

Fig. 2 illustrates a different arrangement of the fluorophores. Again, three oligonucleotides are used. The central oligonucleotide is labelled with a donor fluorophore, 5'-carboxyfluorescein (fluorescein) only. The two adjacent oligonucleotides are each labelled with an individual acceptor fluorophore, 5'-carboxytetramethyl-rhodamine (TMR) and 5'-carboxyrhodamine-X (ROX). These fluorophores have absorption spectra which overlap with fluorescein's emission spectrum and have well separated emission maxima.

On binding of all three oligonucleotides, the fluorophores are positioned such that excitation of fluorescein results in energy transfer from fluorescein to both TMR and ROX. This transfer can again be monitored by observing the emission of the two acceptor fluorophores. Since the acceptors emit at two distinct wavelengths, introduction of a single base mismatch should be detectable by a loss in either or both of these signals.

The positioning of the fluorophores on each oligonucleotide required to optimise energy transfer can be easily determined by the skilled person by preliminary studies. The fluorophores will typically be separated, e.g. by a distance of 7 bases.

FRET assays can be carried out using a fluorimager, irradiating the assays with a laser at a suitable

\$00 mg.

wavelength, e.g. 488 nm for fluorescein, and scanning the emissions using suitable filters, e.g. 530 nm for fluorescein, 570 nm for TMR and 610 nm for ROX.

A characteristic of the common, donor fluorophore is that it should emit at a wavelength that is capable of exciting each of the acceptor fluorophores. The emission of each acceptor fluorophore must be resolvable.

An example of a common fluorophore that can be used as a donor is carboxyfluorescein. Additional examples of suitable acceptors include N,N,N',N'-tetramethyl-6-carboxyrhodamine and 2',7'-dimethoxy-4',5'-dichloro-6-carboxyfluorescein.

The probes may be of DNA. Alternatively, and on the same principle, fluorescently-labelled DNA mimics, e.g. PNA, phosphorothicate DNA, could be used as the probes.

By the use of additional fluorophores with distinct emission profiles, it may be possible to extend this system for the screening of longer target sequences using multiple short complementary strands. This arrangement may permit screening of a section of the DNA target gene in a single homogeneous assay by excitation with, e.g. wavelength laser, or a fluorescence spectrometer, and emission spectrum generating an instantly characterises where, if anywhere, there is a point mutation It is envisaged that sequences of diagnostic or damage. oligonucleotides may be chosen to generate diagnostic kits for screening regions of mutational "hotspots".

The following Example illustrates the invention. In particular, it shows the ability to detect and locate point mutations in a target oligonucleotide.

Example

. :

5

10

15

20

25

30

35

Four target oligonucleotides (SEQ ID Nos 1-4) were designed, each differing only in one nucleotide.

- SEQ ID No 1 5'CGTTCTAAGGATTACGTCGAACCTTTG3'
- SEQ ID No 2 5'CGTTCAAAGGATTACGTCGAACCTTTG3'
 - SEQ ID No 3 5'CGTTCTAAGGATAACGTCGAACCTTTG3'
 - SEQ ID No 4 5'CGTTCTAAGGATTACGTCGAACCATTG3'

5

Sec. Care

Using oligo SEQ ID No 1 as the control, three oligonucleotide probes (SEQ ID No. 5-7) were designed, each capable of hybridising to a distinct region on the control.

SEO ID No 5 3'GCAAGATTC5'

5

10

15

20

25

SEQ ID No 6 3'CTAATGCAG5'

SEQ ID No 7 3'CTTGGAAAC5'

The probes were labelled with a fluorophore at the positions marked (*) as follows: probe SEQ ID No 5 with ROX; probe SEQ ID No 6 with fluorescein; and probe SEQ ID No 7 with TMR.

The emission intensity for both ROX and TMR-labelled probes increased significantly (relative to each singly-bound probe) on hybridisation to the control. However, as shown in Fig. 3, on reaction with each of the target oligonucleotides SEQ ID Nos 2-4, the emission intensity varied significantly, depending on the position of the single nucleotide difference from the control.

This assay provides a simple method for the analysis of hybridisation events in solution and enables detection of mutations within defined regions of a target strand from a single fluorescence measurement. The assay may be modified to utilise immobilised target or probes.

6

SEQUENCE LISTING

(1) GENE	RAL, INFORMATION:	
`_`	APPLICANT: (A) NAME: Cambridge University Technical Services Ltd. (B) STREET: The Old Schools, Trinity Lane (C) CITY: Cambridge (D) STATE: N/A (E) COUNTRY: United Kingdom (F) POSTAL CODE (ZIP): CB2 1TS	
(ii)	TITLE OF INVENTION: DNA MUTATION MAPPING BY MULTIPLE ENERGY TRANSFER INTERACTIONS	
(iii)	NUMBER OF SEQUENCES: 7	
(iv)	COMPUTER READABLE FORM: (A) MEDIUM TYPE: Floppy disk (B) COMPUTER: IBM PC compatible (C) OPERATING SYSTEM: PC-DOS/MS-DOS (D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)	
(v)	CURRENT APPLICATION DATA: APPLICATION NUMBER: WO Not yet known	
(2) INFO	RMATION FOR SEQ ID NO: 1:	
(i)	SEQUENCE CHARACTERISTICS: (A) LENGTH: 27 base pairs (B) TYPE: nucleic acid (C) STRANDEDNESS: single (D) TOPOLOGY: linear	
(ii)	MOLECULE TYPE: other nucleic acid (A) DESCRIPTION: /desc = "Oligonucleotide"	
(xi)	SEQUENCE DESCRIPTION: SEQ ID NO: 1:	
CGTTCTAA	AGG ATTACGTCGA ACCTTTG	27
(2) INFO	RMATION FOR SEQ ID NO: 2:	
(i)	SEQUENCE CHARACTERISTICS: (A) LENGTH: 27 base pairs (B) TYPE: nucleic acid (C) STRANDEDNESS: single (D) TOPOLOGY: linear	

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

CGTTCAAAGG ATTACGTCGA ACCTTTG

WO 98/48048	PCT/GB98/01138
WU 90/40040	FC1/GB96/01136

******* ****_

9.

. :	7	
	(2) INFORMATION FOR SEQ ID NO: 3:	
	(i) SEQUENCE CHARACTERISTICS: (A) LENGTH: 27 base pairs (B) TYPE: nucleic acid (C) STRANDEDNESS: single (D) TOPOLOGY: linear	
	<pre>(ii) MOLECULE TYPE: other nucleic acid (A) DESCRIPTION: /desc = "Oligonucleotide"</pre>	
	(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:	
	CGTTCTAAGG ATAACGTCGA ACCATTG	27
	(2) INFORMATION FOR SEQ ID NO: 4:	
	 (i) SEQUENCE CHARACTERISTICS: (A) LENGTH: 27 base pairs (B) TYPE: nucleic acid (C) STRANDEDNESS: single (D) TOPOLOGY: linear 	
	<pre>(ii) MOLECULE TYPE: other nucleic acid (A) DESCRIPTION: /desc = "Oligonucleotide"</pre>	
	(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:	
	CGTTCTAAGG ATTACGTCGA ACCATTG	27
	(2) INFORMATION FOR SEQ ID NO: 5:	
	 (i) SEQUENCE CHARACTERISTICS: (A) LENGTH: 9 base pairs (B) TYPE: nucleic acid (C) STRANDEDNESS: single (D) TOPOLOGY: linear 	
	<pre>(ii) MOLECULE TYPE: other nucleic acid (A) DESCRIPTION: /desc = "Oligonucleotide"</pre>	

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

GCAAGATTC

PCT/GB98/01138 WO 98/48048

2) INFORMATION	FOR	SEQ	ID	NO:	6:
----------------	-----	-----	----	-----	----

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 9 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: other nucleic acid
 - (A) DESCRIPTION: /desc = "Oligonucleotide"
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

CTAATGCAG

\$00 mm

- (2) INFORMATION FOR SEQ ID NO: 7:
 - (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 9 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

 - (ii) MOLECULE TYPE: other nucleic acid
 (A) DESCRIPTION: /desc = "Oligonucleotide"
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

CTTGGAAAC

9

CLAIMS

10

15

30

- A method for determining the presence and location of a mismatch in a target sequence, which comprises contacting sequence with first, the target second and oligonucleotides capable of hybridising to the natural sequence, in juxtaposition, wherein the oligonucleotides are respectively labelled with first, second and third markers capable of absorption at first, second and third wavelengths and emission at first, second and third wavelengths, such that there is resonance between either or each of first and second markers and between second and third markers, and observing the presence or absence of resonance.
 - 2. A method according to claim 1, wherein the second oligonucleotide comprises two second markers resonating with the first and third markers respectively.
 - 3. A method according to claim 1 or claim 2, wherein one or more markers is a fluorescent moiety.
- 4. A method according to any preceding claim, wherein the one or more markers are selected from 5'-carboxyrhodamine-X, 5'-carboxytetramethylrhodamine, N,N,N',N'-tetramethyl-6-carboxyrhodamine, 2',7'-dimethoxy-4',5'-dichloro-6-carboxyfluorescein and 5'-carboxyfluorescein.
- 5. A method according to any preceding claim, wherein the oligonucleotides each comprise 8-13 bases.
 - 6. A method according to any preceding claim, wherein adjacent markers are separated by at least 7 bases.
 - 7. A method according to any preceding claim, wherein the target or the first, second and third oligonucleotides are immobilised.
 - 8. A kit comprising, in separate compartments, first, second and third markers defined in any of claims 1 to 5.

1/2

FIG. 1




FIG. 2





FIG. 3



PCT


WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)_

(51) International Patent Classification ⁶ : C12Q 1/68	А3	(11) International Publication Number: WO 98/4 (43) International Publication Date: 29 October 1998 (29)	
(21) International Application Number: PCT/GB (22) International Filing Date: 20 April 1998 ((30) Priority Data: 9707996.6 21 April 1997 (21.04.97) (71) Applicant (for all designated States except US): CAM UNIVERSITY TECHNICAL SERVICES LTD. [The Old Schools, Trinity Lane, Cambridge CB2]	20.04.9 ((BRIDG [GB/GI	BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, G GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, K LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, S TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), E patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), Eu patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, G) IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, E CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	GB, GE, KR, KZ, N, MW, SK, SL, ARIPO Eurasian uropean BB, GR, BJ, CF,
 (72) Inventor; and (75) Inventor/Applicant (for US only): BALASUBRAM Shankar [GB/GB]; University of Cambridge, I Chemistry, Lensfield Road, Cambridge CB2 1EW (74) Agent: GILL JENNINGS & EVERY; Broadgate Eldon Street, London EC2M 7LH (GB). 	Dept. (GB).	of (88) Date of publication of the international search report: 21 January 1999 (2)	

(54) Title: DNA MUTATION MAPPING BY MULTIPLE ENERGY TRANSFER INTERACTIONS



(57) Abstract

A method for determining the presence and location of a mismatch in a target sequence using Fluorescence Resonance Energy Transfer (FRET). The method comprises contacting the target sequence with at least three labelled oligonucleotide probes capable of hybridising to the natural sequence, in juxtaposition. The central probe is labelled with a donor fluorophore and the probe either side of this is labelled with a distinct acceptor fluorophore. Hybridisation to the target sequence results in resonance between the donor fluorophore and the acceptor fluorophores. A mismatch present in the target sequence will disrupt hybridisation to that region, resulting in a significant alteration to the resonance signal.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

	-	-	-				
AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
ΑT	Austria	FR	France	LU	Luxembourg	SN	Senegal
ΑU	Australia .	GA	Gabon	LV	Latvia	SZ	Swaziland
ΑZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghann	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	1E	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	n.	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MIX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JР	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	zw	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
cu	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

INTERNATIONAL SEARCH REPORT

Inte. .onal Application No PCT/GB 98/01138

A. CLASSIF IPC 6	C 12Q1/68		:
According to	International Patent Classification (IPC) or to both national classification	ation and IPC	
	SEARCHED		
Minimum do IPC 6	cumentation searched (classification system followed by classification $C12Q$	on symbols)	
Documentati	ion searched other than minimum documentation to the extent that si	uch documents are included in the field	is searched
Electronic da	ata base consulted during the international search (name of data ba	se and, where practical, search terms (used)
C. DOCUME	ENTS CONSIDERED TO BE RELEVANT		
Category :	Citation of document, with indication, where appropriate, of the rek	evant passages	Relevant to claim No.
X	WO 96 25518 A (SOCIETY FOR TECHNO INNOVATION ;MASUKO MASAYUKI (JP) KATSUYO) 22 August 1996 see figure 3 & EP 0 810 291 A see column 3, line 41 - line 47;	; EBATA	1-8
X	EP 0 229 943 A (MOLECULAR BIOSYS 29 July 1987 see page 13, last paragraph; fig		1-8
A	WO 93 09128 A (NANOTRONICS INC) 13 May 1993		
		-/	
X Furt	ther documents are listed in the continuation of box C.	Patent family members are I	isted in annex.
"A" docum	ategories of cited documents : ent defining the general state of the art which is not dered to be of particular relevance	"T" later document published after th or priority date and not in conflic cited to understand the principle invention	t with the application but
"L" docume which	document but published on or after the international date ent which may throw doubte on priority daim(s) or is cited to establish the publication date of another in or other special reason (as specified)	"X" document of particular relevance cannot be considered novel or of involve an invertilve step when it "Y" document of particular relevance cannot be considered to involve	annot be considered to the document is taken alone ; the claimed invention
other "P" docum	nent referring to an oral disclosure, use, exhibition or means ent published prior to the international filing date but than the priority date daimed	document is combined with one ments, such combination being in the art. "8" document member of the same p	or more other such docu- obvious to a person skilled
<u> </u>	actual completion of theinternational search	Date of mailing of the internation	
1	16 October 1998	26/10/1998	
Name and	maiting address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk	Authorized officer	
	Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016	Molina Galan,	E

1

INTERNATIONAL SEARCH REPORT

Ints Jonal Application No PCT/GB 98/01138

	·	PCT/GB 98/01138						
.(Continu	ontinuation) DOCUMENTS CONSIDERED TO BE RELEVANT							
ategory ·	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.						
A	TYAGI S ET AL: "MOLECULAR BEACONS: PROBES THAT FLUORESCE UPON HYBRIDIZATION" BIO/TECHNOLOGY, vol. 14, 1 March 1996, pages 303-308, XP000196024 cited in the application							
A	EP 0 601 889 A (MAINE MEDICAL CENTER RES) 15 June 1994							
	·	·						
•								

INTERNATIONAL SEARCH REPORT

Information on patent family members

Inte Jonal Application No
PCT/GB 98/01138

Patent document cited in search report	:	Publication date		atent family member(s)	Publication date
WO 9625518	Α	22-08-1996	AU	694313 B	16-07-1998
			AU	4633896 A	04-09-1996
			CA	2213240 A	22-08-1996
			EP	0810291 A	03-12-1997
EP 0229943	Α	29-07-1987	CA	1273552 A	04-09-1990
			DE	3681272 A	10-10-1991
			GR	3002735 T	25-01-1993
			JP	2033069 C	19-03-1996
			JP	7063400 B	12-07-1995
			JP	62157570 A	13-07-1987
			US	4996143 A	26-02-1991
WO 9309128	Α	13-05-1993	AU	667497 B	28-03-1996
			AU	3136493 A	07-06-1993
			CA	2123133 A	13-05-1993
			EP	0620822 A	26-10-1994
			JP	7502992 T	30-03-1995
			US	5565322 A	15-10-1996
			US	5532129 A	02-07-1996
			US	5787032 A	28-07-1998
EP 0601889	Α	15-06-1994	US	5607834 A	04-03-1997